

Metode iterasi Gauss-Seidel

Algoritma

A set of n equations and n unknowns:

$$a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \dots + a_{1n}x_n = b_1$$

$$a_{21}x_1 + a_{22}x_2 + a_{23}x_3 + \dots + a_{2n}x_n = b_2$$

$$a_{n1}x_1 + a_{n2}x_2 + a_{n3}x_3 + \dots + a_{nn}x_n = b_n$$

If: the diagonal elements are non-zero

Rewrite each equation solving for the corresponding unknown

ex:

First equation, solve for x_1

Second equation, solve for x_2

Gauss-Seidel Method

Algorithm

Rewriting each equation

$$x_1 = \frac{c_1 - a_{12}x_2 - a_{13}x_3 - \dots - a_{1n}x_n}{a_{11}} \quad \leftarrow \text{From Equation 1}$$

$$x_2 = \frac{c_2 - a_{21}x_1 - a_{23}x_3 - \dots - a_{2n}x_n}{a_{22}} \quad \leftarrow \text{From equation 2}$$
$$\vdots \quad \vdots \quad \vdots$$

$$x_{n-1} = \frac{c_{n-1} - a_{n-1,1}x_1 - a_{n-1,2}x_2 - \dots - a_{n-1,n-2}x_{n-2} - a_{n-1,n}x_n}{a_{n-1,n-1}} \quad \leftarrow \text{From equation n-1}$$

$$x_n = \frac{c_n - a_{n1}x_1 - a_{n2}x_2 - \dots - a_{n,n-1}x_{n-1}}{a_{nn}} \quad \leftarrow \text{From equation n}$$

Gauss-Seidel Method

Algorithm

General Form of each equation

$$x_1 = \frac{c_1 - \sum_{\substack{j=1 \\ j \neq 1}}^n a_{1j} x_j}{a_{11}}$$
$$x_{n-1} = \frac{c_{n-1} - \sum_{\substack{j=1 \\ j \neq n-1}}^n a_{n-1,j} x_j}{a_{n-1,n-1}}$$

$$x_2 = \frac{c_2 - \sum_{\substack{j=1 \\ j \neq 2}}^n a_{2j} x_j}{a_{22}}$$
$$x_n = \frac{c_n - \sum_{\substack{j=1 \\ j \neq n}}^n a_{nj} x_j}{a_{nn}}$$

Gauss-Seidel Method

Solve for the unknowns

Assume an initial guess for [X]

$$\begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_{n-1} \\ x_n \end{bmatrix}$$

Use rewritten equations to solve for each value of x_i .

Important: Remember to use the most recent value of x_i . Which means to apply values calculated to the calculations remaining in the **current** iteration.

Gauss-Seidel Method

Calculate the Absolute Relative Approximate Error

$$|E_a|_i = \left| \frac{x_i^{new} - x_i^{old}}{x_i^{new}} \right| \times 100$$

So when has the answer been found?

The iterations are stopped when the absolute relative approximate error is less than a prespecified tolerance for all unknowns.

Example

Given the system of equations

$$12x_1 + 3x_2 - 5x_3 = 1$$

$$x_1 + 5x_2 + 3x_3 = 28$$

$$3x_1 + 7x_2 + 13x_3 = 76$$

With an initial guess of

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$$

The coefficient matrix is:

$$[A] = \begin{bmatrix} 12 & 3 & -5 \\ 1 & 5 & 3 \\ 3 & 7 & 13 \end{bmatrix}$$

Will the solution converge using the Gauss-Siedel method?

Checking if the coefficient matrix is diagonally dominant

$$[A] = \begin{bmatrix} 12 & 3 & -5 \\ 1 & 5 & 3 \\ 3 & 7 & 13 \end{bmatrix}$$
$$|a_{11}| = |12| = 12 \geq |a_{12}| + |a_{13}| = |3| + |-5| = 8$$
$$|a_{22}| = |5| = 5 \geq |a_{21}| + |a_{23}| = |1| + |3| = 4$$
$$|a_{33}| = |13| = 13 \geq |a_{31}| + |a_{32}| = |3| + |7| = 10$$

Rewriting each equation

$$\begin{bmatrix} 12 & 3 & -5 \\ 1 & 5 & 3 \\ 3 & 7 & 13 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} 1 \\ 28 \\ 76 \end{bmatrix}$$

$$x_1 = \frac{1 - 3x_2 + 5x_3}{12}$$

$$x_2 = \frac{28 - x_1 - 3x_3}{5}$$

$$x_3 = \frac{76 - 3x_1 - 7x_2}{13}$$

With an initial guess of

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$$

$$x_1 = \frac{1 - 3(0) + 5(1)}{12} = 0.50000$$

$$x_2 = \frac{28 - (0.5) - 3(1)}{5} = 4.9000$$

$$x_3 = \frac{76 - 3(0.50000) - 7(4.9000)}{13} = 3.0923$$

| Iteration | a_1 | $ \mathcal{E}_a _1$ | a_2 | $ \mathcal{E}_a _2$ | a_3 | $ \mathcal{E}_a _3$ |
|-----------|---------|---------------------|--------|---------------------|--------|---------------------|
| 1 | 0.50000 | 67.662 | 4.900 | 100.00 | 3.0923 | 67.662 |
| 2 | 0.14679 | 240.62 | 3.7153 | 31.887 | 3.8118 | 18.876 |
| 3 | 0.74275 | 80.23 | 3.1644 | 17.409 | 3.9708 | 4.0042 |
| 4 | 0.94675 | 21.547 | 3.0281 | 4.5012 | 3.9971 | 0.65798 |
| 5 | 0.99177 | 4.5394 | 3.0034 | 0.82240 | 4.0001 | 0.07499 |
| 6 | 0.99919 | 0.74260 | 3.0001 | 0.11000 | 4.0001 | 0.00000 |

The solution obtained

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0.99919 \\ 3.0001 \\ 4.0001 \end{bmatrix}$$

is close to the exact solution of

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 1 \\ 3 \\ 4 \end{bmatrix}$$