



Gauss Quadrature Rule of Integration

Nur Laila Hamidah

Capaian Pembelajaran

- Mahasiswa mampu menerapkan **metode numerik** Gauss Quadrature Rule untuk penyelesaian persamaan **integral tertentu**

Gauss Quadrature Rule of Integration

[http://numericalmethods.eng.usf.
edu](http://numericalmethods.eng.usf.edu)

What is Integration?

Integration

The process of measuring the area under a curve.


$$I = \int_a^b f(x) dx$$

Where:

$f(x)$ is the integrand

a = lower limit of integration

b = upper limit of integration



TWO-POINT GAUSSIAN QUADRATURE RULE

Basis of the Gaussian Quadrature Rule

Previously, the Trapezoidal Rule was developed by the method of undetermined coefficients. The result of that development is summarized below.

$$\int_a^b f(x)dx \approx c_1 f(a) + c_2 f(b)$$
$$= \frac{b-a}{2} f(a) + \frac{b-a}{2} f(b)$$

Basis of the Gaussian Quadrature Rule

The two-point Gauss Quadrature Rule is an extension of the Trapezoidal Rule approximation where the arguments of the function are not predetermined as a and b but as unknowns x_1 and x_2 . In the two-point Gauss Quadrature Rule, the integral is approximated as

$$I = \int_a^b f(x) dx \approx c_1 f(x_1) + c_2 f(x_2)$$

Basis of the Gaussian Quadrature Rule

The four unknowns x_1, x_2, c_1 and c_2 are found by assuming that the formula gives exact results for integrating a general third order polynomial, $f(x) = a_0 + a_1x + a_2x^2 + a_3x^3$.

Hence

$$\begin{aligned}\int_a^b f(x) dx &= \int_a^b (a_0 + a_1x + a_2x^2 + a_3x^3) dx \\ &= \left[a_0x + a_1 \frac{x^2}{2} + a_2 \frac{x^3}{3} + a_3 \frac{x^4}{4} \right]_a^b \\ &= a_0(b-a) + a_1 \left(\frac{b^2 - a^2}{2} \right) + a_2 \left(\frac{b^3 - a^3}{3} \right) + a_3 \left(\frac{b^4 - a^4}{4} \right)\end{aligned}$$

Basis of the Gaussian Quadrature Rule

It follows that

$$\int_a^b f(x) dx = c_1(a_0 + a_1x_1 + a_2x_1^2 + a_3x_1^3) + c_2(a_0 + a_1x_2 + a_2x_2^2 + a_3x_2^3)$$

Equating Equations the two previous two expressions yield

$$\begin{aligned} & a_0(b-a) + a_1\left(\frac{b^2 - a^2}{2}\right) + a_2\left(\frac{b^3 - a^3}{3}\right) + a_3\left(\frac{b^4 - a^4}{4}\right) \\ &= c_1(a_0 + a_1x_1 + a_2x_1^2 + a_3x_1^3) + c_2(a_0 + a_1x_2 + a_2x_2^2 + a_3x_2^3) \\ &= a_0(c_1 + c_2) + a_1(c_1x_1 + c_2x_2) + a_2(c_1x_1^2 + c_2x_2^2) + a_3(c_1x_1^3 + c_2x_2^3) \end{aligned}$$

Basis of the Gaussian Quadrature Rule

Since the constants a_0, a_1, a_2, a_3 are arbitrary

$$b - a = c_1 + c_2$$

$$\frac{b^2 - a^2}{2} = c_1 x_1 + c_2 x_2$$

$$\frac{b^3 - a^3}{3} = c_1 x_1^2 + c_2 x_2^2$$

$$\frac{b^4 - a^4}{4} = c_1 x_1^3 + c_2 x_2^3$$

Basis of Gauss Quadrature

The previous four simultaneous nonlinear Equations have only one acceptable solution,

$$x_1 = \left(\frac{b-a}{2}\right)\left(-\frac{1}{\sqrt{3}}\right) + \frac{b+a}{2}$$

$$x_2 = \left(\frac{b-a}{2}\right)\left(\frac{1}{\sqrt{3}}\right) + \frac{b+a}{2}$$

$$c_1 = \frac{b-a}{2}$$

$$c_2 = \frac{b-a}{2}$$

Basis of Gauss Quadrature

Hence Two-Point Gaussian Quadrature Rule

$$\int_a^b f(x)dx \approx c_1 f(x_1) + c_2 f(x_2)$$
$$= \frac{b-a}{2} f\left(\frac{b-a}{2}\left(-\frac{1}{\sqrt{3}}\right) + \frac{b+a}{2}\right) + \frac{b-a}{2} f\left(\frac{b-a}{2}\left(\frac{1}{\sqrt{3}}\right) + \frac{b+a}{2}\right)$$

HIGHER POINT GAUSSIAN QUADRATURE FORMULAS

Higher Point Gaussian Quadrature Formulas

$$\int_a^b f(x)dx \approx c_1 f(x_1) + c_2 f(x_2) + c_3 f(x_3)$$

is called the three-point Gauss Quadrature Rule.

The coefficients c_1 , c_2 , and c_3 , and the functional arguments x_1 , x_2 , and x_3 are calculated by assuming the formula gives exact expressions for integrating a fifth order polynomial

$$\int_a^b (a_0 + a_1 x + a_2 x^2 + a_3 x^3 + a_4 x^4 + a_5 x^5) dx$$

General n-point rules would approximate the integral

$$\int_a^b f(x)dx \approx c_1 f(x_1) + c_2 f(x_2) + \dots + c_n f(x_n)$$

Arguments and Weighing Factors for n-point Gauss Quadrature Formulas

In handbooks, coefficients and arguments given for n-point Gauss Quadrature Rule are given for integrals

$$\int_{-1}^1 g(x) dx \cong \sum_{i=1}^n c_i g(x_i)$$

as shown in Table 1.

Table 1: Weighting factors c and function arguments x used in Gauss Quadrature Formulas.

Points	Weighting Factors	Function Arguments
2	$c_1 = 1.000000000$ $c_2 = 1.000000000$	$x_1 = -0.577350269$ $x_2 = 0.577350269$
3	$c_1 = 0.555555556$ $c_2 = 0.888888889$ $c_3 = 0.555555556$	$x_1 = -0.774596669$ $x_2 = 0.000000000$ $x_3 = 0.774596669$
4	$c_1 = 0.347854845$ $c_2 = 0.652145155$ $c_3 = 0.652145155$ $c_4 = 0.347854845$	$x_1 = -0.861136312$ $x_2 = -0.339981044$ $x_3 = 0.339981044$ $x_4 = 0.861136312$

Arguments and Weighing Factors for n-point Gauss Quadrature Formulas

Table 1 (cont.) : Weighting factors c and function arguments x used in Gauss Quadrature Formulas.

Points	Weighting Factors	Function Arguments
5	$c_1 = 0.236926885$ $c_2 = 0.478628670$ $c_3 = 0.568888889$ $c_4 = 0.478628670$ $c_5 = 0.236926885$	$x_1 = -0.906179846$ $x_2 = -0.538469310$ $x_3 = 0.000000000$ $x_4 = 0.538469310$ $x_5 = 0.906179846$
6	$c_1 = 0.171324492$ $c_2 = 0.360761573$ $c_3 = 0.467913935$ $c_4 = 0.467913935$ $c_5 = 0.360761573$ $c_6 = 0.171324492$	$x_1 = -0.932469514$ $x_2 = -0.661209386$ $x_3 = -0.2386191860$ $x_4 = 0.2386191860$ $x_5 = 0.661209386$ $x_6 = 0.932469514$

Arguments and Weighing Factors for n -point Gauss Quadrature Formulas

So if the table is given for $\int_{-1}^1 g(x) dx$ integrals, how does one solve $\int_a^b f(x) dx$? The answer lies in that any integral with limits of $[a, b]$ can be converted into an integral with limits $[-1, 1]$ Let

$$x = mt + c$$

$$\text{If } x = a, \text{ then } t = -1$$

$$\text{If } x = b, \text{ then } t = 1$$

Such that:

$$m = \frac{b - a}{2}$$

Arguments and Weighing Factors for n -point Gauss Quadrature Formulas

Then $c = \frac{b+a}{2}$ Hence

$$x = \frac{b-a}{2}t + \frac{b+a}{2} \quad dx = \frac{b-a}{2}dt$$

Substituting our values of x , and dx into the integral gives us

$$\int_a^b f(x)dx = \int_{-1}^1 f\left(\frac{b-a}{2}t + \frac{b+a}{2}\right) \frac{b-a}{2} dt$$

Example 1

For an integral $\int_a^b f(x) dx$, derive the one-point Gaussian Quadrature Rule.

Solution

The one-point Gaussian Quadrature Rule is

$$\int_a^b f(x) dx \approx c_1 f(x_1)$$

Solution

The two unknowns x_1 , and c_1 are found by assuming that the formula gives exact results for integrating a general first order polynomial,

$$f(x) = a_0 + a_1x.$$

$$\begin{aligned}\int_a^b f(x)dx &= \int_a^b (a_0 + a_1x)dx \\ &= \left[a_0x + a_1 \frac{x^2}{2} \right]_a^b \\ &= a_0(b - a) + a_1 \left(\frac{b^2 - a^2}{2} \right)\end{aligned}$$

Solution

It follows that

$$\int_a^b f(x)dx = c_1(a_0 + a_1x_1)$$

Equating Equations, the two previous two expressions yield

$$a_0(b - a) + a_1\left(\frac{b^2 - a^2}{2}\right) = c_1(a_0 + a_1x_1) = a_0(c_1) + a_1(c_1x_1)$$

Basis of the Gaussian Quadrature Rule

Since the constants a_0 , and a_1 are arbitrary

$$b - a = c_1$$

$$\frac{b^2 - a^2}{2} = c_1 x_1$$

giving

$$c_1 = b - a$$

$$x_1 = \frac{b + a}{2}$$

Solution

Hence One-Point Gaussian Quadrature Rule

$$\int_a^b f(x)dx \approx c_1 f(x_1) = (b-a) f\left(\frac{b+a}{2}\right)$$

Example 2

- a) Use two-point Gauss Quadrature Rule to approximate the distance covered by a rocket from $t=8$ to $t=30$ as given by

$$x = \int_8^{30} \left(2000 \ln \left[\frac{140000}{140000 - 2100t} \right] - 9.8t \right) dt$$

- b) Find the true error, E_t for part (a).
- c) Also, find the absolute relative true error, $|\epsilon_a|$ for part (a).

Solution

First, change the limits of integration from $[8,30]$ to $[-1,1]$ by previous relations as follows

$$\begin{aligned}\int_8^{30} f(t) dt &= \frac{30-8}{2} \int_{-1}^1 f\left(\frac{30-8}{2}x + \frac{30+8}{2}\right) dx \\ &= 11 \int_{-1}^1 f(11x + 19) dx\end{aligned}$$

Solution (cont)

Next, get weighting factors and function argument values from Table 1 for the two point rule,

$$c_1 = 1.0000000000$$

$$x_1 = -0.577350269$$

$$c_2 = 1.0000000000$$

$$x_2 = 0.577350269$$

Solution (cont.)

Now we can use the Gauss Quadrature formula

$$\begin{aligned} 11 \int_{-1}^1 f(11x + 19) dx &\approx 11c_1 f(11x_1 + 19) + 11c_2 f(11x_2 + 19) \\ &= 11f(11(-0.5773503) + 19) + 11f(11(0.5773503) + 19) \\ &= 11f(12.64915) + 11f(25.35085) \\ &= 11(296.8317) + 11(708.4811) \\ &= 11058.44 \text{ m} \end{aligned}$$

Solution (cont)

since

$$\begin{aligned} f(12.64915) &= 2000 \ln \left[\frac{140000}{140000 - 2100(12.64915)} \right] - 9.8(12.64915) \\ &= 296.8317 \end{aligned}$$

$$\begin{aligned} f(25.35085) &= 2000 \ln \left[\frac{140000}{140000 - 2100(25.35085)} \right] - 9.8(25.35085) \\ &= 708.4811 \end{aligned}$$

Solution (cont)

b) The true error, E_t , is

$$\begin{aligned} E_t &= \text{True Value} - \text{Approximate Value} \\ &= 11061.34 - 11058.44 \\ &= 2.9000 \text{ m} \end{aligned}$$

c) The absolute relative true error, $|\epsilon_t|$, is (Exact value = 11061.34m)

$$|\epsilon_t| = \left| \frac{11061.34 - 11058.44}{11061.34} \right| \times 100\%$$

$$= 0.0262\%$$